arXiv:0710.4574v3 [cond-mat.str-el] 10 Oct 2008

Spontaneous spin ordering of Dirac spin liquid in a magnetic field

Ying Ran,[1] Wing-Ho Ko,[2] Patrick A. Lee,[2] and Xiao-Gang Wen[2]
[1]*Department of Physics, University of California, Berkeley, California 94720*
[2]*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*
(Dated: October 10, 2008)

The Dirac spin liquid was proposed to be the ground state of the spin-1/2 Kagome antiferromagnets. In a magnetic field B, we show that the state with Fermi pocket is unstable to the Landau level (LL) state. The LL state breaks the spin rotation around the axis of the magnetic field. We find that the LL state has an in-plane 120° $q = 0$ magnetization M which scales with the external field $M \sim B^\alpha$, where α is an intrinsic calculable universal number of the Dirac spin liquid. We discuss the related experimental implications which can be used to detect the possible Dirac spin liquid phase in Herbertsmithite $\mathrm{ZnCu_3(OH)_6Cl_2}$.

PACS numbers: 75.10.Jm, 75.50.Ee

Spin liquids (SL), defined as the ground states of spin systems with half integer spins per unit cell which does not order magnetically and/or break translation symmetry, are believed to contain fundamentally new physics beyond the Landau's symmetry breaking characterization of phases. After years of search, a promising candidate finally emerged in the spin-1/2 Kagome system Herbertsmithite $\mathrm{ZnCu_3(OH)_6Cl_2}$[1, 2, 3]. Despite an antiferromagnetic exchange $J \approx 170 \sim 190$K[4, 5], the system does not order down to 50mK[1, 2, 3]. Theoretically the Dirac-SL is proposed[6] as the ground state of the nearest neighbor Heisenberg antiferromagnetic model on Kagome lattice. Unfortunately the spin susceptibility χ_s is consistent with $\sim 4\%$ magnetic impurities[4], and possible Dzyaloshinskii-Moriya interaction has also been proposed[5]. These may explain the hump of the specific heat C around 2K and also obscure the $C \propto T^2$ and $\chi_s \propto T$ behaviors predicted by the Dirac-SL. In this paper we propose a unique signature of the Dirac-SL in an external magnetic field, which can be used to detect the Dirac-SL phase in experiment and numerical simulations.

The Dirac-SL experiences no orbital effect in the external magnetic field because it is an insulator. The Zeeman coupling $g\mu_B \vec{B}\cdot\sum_i \vec{S}_i$ polarizes spin along the field direction, breaks time reversal and breaks $SU(2)$ spin rotation down to $U(1)$. Let us denote the direction of $\vec{B}$ as z-direction (note that the xy-plane does not have to be the plane of the two-dimensional Kagome system). We find that the Dirac-SL will spontaneously break the remaining S_z-$U(1)$ and form a staggered magnetization M (see Fig.2(b)) in the xy-plane which scales with B: $M \sim B^\alpha$. The positive exponent α is an intrinsic universal number of the Dirac-SL phase and in principle calculable.

This unique signature of the Dirac-SL can be compared with a regular co-planer antiferromagnetic (AF) ordered phase. In a small magnetic field B, the magnetization of the regular AF phase would rotate into the xy-plane to maximize the susceptibility along the B direction. As B is tuned to zero, the in-plane magnetization remains finite. In contrast, for the Dirac-SL we predict that M vanishes as B^α. This suggests that the Dirac-SL can be viewed as an AF phase whose long-range AF order has been destroyed by the quantum fluctuations.

We begin with writing down the the low energy theory of the Dirac-SL[6], which includes four flavors of fermions coupled with compact $U(1)$ gauge field in 2+1 dimension (QED$_3$):

$$S = \int dx^3 \Big[\frac{1}{g^2}(\varepsilon_{\lambda\mu\nu}\partial_\mu a_\nu)^2 + \sum_\sigma \bar{\psi}_{+\sigma}\left(\partial_\mu - i a_\mu\right)\tau_\mu \psi_{+\sigma} + \sum_\sigma \bar{\psi}_{-\sigma}\left(\partial_\mu - i a_\mu\right)\tau_\mu \psi_{-\sigma}\Big] + \cdots , \tag{1}$$

where the two-component fermionic Dirac spinon fields are denoted by $\psi_{\pm\sigma}$, where $\pm$ label the two inequivalent nodes and σ the up/down spins.

The spinon carries spin-1/2 and charge-0, and couples to the external magnetic field only by the Zeeman effect. In the presence of a magnetic field, the simplest guess is that the Dirac points will change into Fermi pockets due to Zeeman splitting. We will call this state Fermi pocket state (FP). But, can there be other states whose energy may be lower?

The FP state has many gapless spinon excitations near the Fermi pockets and in general not energetically favorable. One natural way to gap out the Fermi pocket state is to induce (internal) gauge fluxes and develop Landau levels. Here we take advantage of the appearance of zero energy Landau levels when Dirac particles are subject to a magnetic field, as is well known in the recent studies of graphene[7, 8]. If the gauge flux is adjusted in such a way that the zero-energy Landau level is fully filled for up-spin, and fully empty for down-spin, then the spinons are fully gapped (Fig.1). We will call this state Landau level state(LL).

In the following we show that LL state has lower energy than FP state. We first compare the energies of LL state and FP state at the mean-field level (ignoring the gauge fluctuation) with fixed S_z polarization. We set aside the Zeeman energy which is common between the two states.



FIG. 1: The The spinons is gapless in FP state with an electron-like Fermi pocket for spin up spinons and hole-like pocket for spin down spinons. The spinons are fully gapped in the LL state. Each Landau level is doubly degenerate due to two inequivalent Dirac nodes.

The density of spin imbalance is:

$$\Delta n = \frac{\Delta N}{A} = \frac{N_\uparrow - N_\downarrow}{A} = 4 \cdot \frac{1}{4\pi} \left(\frac{\mu_B B}{v_F} \right)^2, \tag{2}$$

and the mean-field energy density is found to be

$$\Delta e^{FP}_{MF} = \frac{\Delta E^{FP}_{MF}}{A} = \frac{2\sqrt{\pi} v_F}{3} (\Delta n)^{\frac{3}{2}}, \tag{3}$$

where $N_\uparrow$ and $N_\downarrow$ are the number of up and down spins, ΔE is the energy increase compared to Dirac point state, A is the system area, B is the magnetic field, and v_F is the mean-field Fermi velocity. The factor 4 in Eq.(2) is from the fact that there are two spin degeneracy and two nodal degeneracy. We choose our units such that $\hbar = 1$.

To make sure the LL state have the same spin imbalance as the FP state, each Landau level should contain $\Delta n A/2$ states, i.e. the induced gauge magnetic field b satisfies $\Delta n = 2 \cdot \frac{b}{2\pi}$, where the 2π is gauge flux quantum. Since at the mean-field level the internal gauge field costs no energy, the energy of the mean-field LL state is just the difference between the sum of the energies of all negative landau levels, and the filled Fermi sea:

$$\begin{aligned} \Delta e^{LL}_{MF} &= 4 v_F \left(-\frac{b}{2\pi} \sum_{n=1}^{\infty} \sqrt{2nb} - \int \frac{dk^2}{(2\pi)^2} (-k) \right) \\ &= \frac{\zeta(\frac{3}{2}) b^{3/2} v_F}{\sqrt{2}\pi^2} = \frac{\zeta(\frac{3}{2}) \Delta n^{3/2} v_F}{\sqrt{2\pi}}. \end{aligned} \tag{4}$$

The same result was obtained in Ref [9, 10]. From Eq.(3,4) we have $\frac{\Delta e^{FP}_{MF}}{\Delta e^{LL}_{MF}} = \frac{2\sqrt{2\pi}}{3\zeta(3/2)} \doteq 1.134$, i.e., the LL state has lower mean field energy than the FP state.

We may also ask whether in graphene a spontaneous magnetic field may be generated by this mechanism, if electrons or holes are introduced by gating. The answer is negative and the difference is that the physical electromagnetic field has an energy cost of $f^2_{\mu\nu}$. Furthermore the field is three dimensional and the magnetic field must form closed lines. The magnetic flux would be divided into domains of size R with opposite signs. The size R can be estimated by minimizing the total energy, which is the sum of the magnetic field energy $\sim B^2 R^3$ and the Landau level energy gain $\sim -v_F \sqrt{\hbar e B/c}\, BR^2/\Phi_0$ where $\Phi_0 = \frac{hc}{e}$ is the flux quantum. The optimal R_* is found to satisfy $\frac{BR_*^2}{\Phi_0} \sim (\alpha \frac{v_F}{c})^2$, where $\alpha = \frac{e^2}{\hbar c} = \frac{1}{137}$ is the fine structure constant. This means that the flux through the entire domain is $\sim 10^{-7} \Phi_0$, which is not self consistent with the condition that there is at least one flux quantum through the domain to support Landau levels.



FIG. 2: (color online) (a) We convert the data in Table I into energy density Δe and spin imbalance density Δn (units defined in text), and plot them together with the fitted scaling form $\Delta e = a \Delta n^{3/2} + b \Delta n^2$. (b): The XY spin order in the LL state. We also present the relative phase of S^+_i operator computed by projected landau level state in unit of π (see Eq.(7)). Note that we fix the left-bottom corner site as the site i in Eq.(7), and scan the site j for the nine sites depicted.

To include the effect of gauge fluctuations and to go beyond the mean field theory, we will use the Gutzwiller projected wavefunction to calculate the energies of LL and FP states. To obtain the Gutzwiller projected wavefunction we first write down the mean-field Hamiltonian on lattice: $H_{mean} = \sum_{ij} \chi_{ij} f^\dagger_{i\sigma} f_{j\sigma}$, where $f_{i\sigma}$ are the fermionic spinons. The mean-field ground state $|\Psi_{mean}(\chi_{ij})\rangle$ with mean-field parameters χ_{ij} is a spin singlet. The projected wavefunction $|\Psi_{prj}(\chi_{ij})\rangle = P|\Psi_{mean}(\chi_{ij})\rangle$ removes the unphysical states and becomes a spin state; here $P = \prod_i (1 - n_{i\uparrow} n_{i\downarrow})$ is the projection operator ensuring one fermion per site. The physical observables can be measured on $|\Psi_{prj}(\chi_{ij})\rangle$ by a Monte Carlo approach[11]. The Dirac-SL is characterized by χ_{ij} such that $|\chi_{ij}| = \chi$ is a bond independent constant and there are π fluxes through the hexagonal plaquettes and 0 fluxes through the triangular plaquettes[6]. Note that the projected Dirac-SL state has no tunable parameter since χ only gives the wavefunction an overall factor. On a 16x16 unit cell lattice, the energetics of the FP and LL states are given in Table I.

Is the $\Delta e \sim \Delta n^{3/2}$ law still valid after projection? The answer is positive because the S_z and energy are conserved and have no anomalous dimension. The energies in Table I can be thus fitted (Fig.2(a)) as

$$\Delta e^{Prj}_{FP} = 0.33(2) \Delta n^{3/2} + 0.00(4) \Delta n^2, \tag{5}$$

$$\Delta e^{Prj}_{LL} = 0.223(6) \Delta n^{3/2} + 0.03(1) \Delta n^2, \tag{6}$$

where the units are chosen such that the unit cell spacing $a = 1$ and $J = 1$ and we include the first order correction to scaling, the Δn^2 term. From the coefficient 0.33(2) in

ΔN	0	8	16	32	40	56
$E^{FP}_{\rm Prj}$	-0.42865(2)	-0.42798(2)	-0.42688(2)	-0.42327(2)	-0.42131(2)	-0.41638(2)
$E^{LL}_{\rm Prj}$	-0.42865(2)	-0.42817(2)	-0.42732(2)	-0.42493(2)	-0.42346(2)	-0.41994(2)

TABLE I: The per-site energies of the projected FP and LL states in unit of Heisenberg coupling J on a 16x16 unit cell lattice with periodic-antiperiodic boundary conditions. ΔN is the spin imbalance. The FP states has no total gauge flux through the sample, while the LL state has $\frac{\Delta N}{2}$ flux quanta through the sample which is uniformly distributed in the triangular and hexagonal plaquettes.

Eq.(5) we are able to read off the effective fermi velocity $v_F^* = \frac{3\cdot 0.33(2)}{2\sqrt{\pi}} = 0.28(2)\frac{Ja}{\hbar}$ by fitting to the free fermion result Eq.(3). v_F^* is almost twice the mean-field value $v_F = \frac{a\chi}{\sqrt{2}\hbar}$ with $\chi = 0.221J$[12]. For Herbertsmithite this means $v_F^* = 4.9\cdot 10^3$m/s assuming $J = 200$K. This is very close to the fermi velocity we found by a projected band structure study [13], where only one particle-hole excitation is considered. The closeness of the two results imply that the gauge interactions between many particle-hole excitations may only give small corrections for energetics.

From Eq. (5,6) we see that the LL state has a lower energy than the FP state. Therefore the LL state may be the true ground state in the presence of a magnetic field. Due the presence of internal gauge flux, one can easily see that the LL state breaks parity (mirror reflection). Surprisingly, it turns out that the LL state breaks the S_z spin rotation as well. To understand this let us consider the low energy collective excitations in the LL state. Since the spinons ψ are gapped due to the Landau levels, the gauge field a_μ is the only low energy excitation. We note that the total Hall conductance of the spin-up spinons is $+1$ and the total Hall conductance for the spin-down spinons is -1. Thus the total Hall conductance is zero and there is no Chern-Simons term for the gauge field a_μ. As a result, the dynamics of the gauge field is controlled by the Maxwell term $f_{\mu\nu}^2$, which arise from integrating out high-energy fermions, and gives rise to a linear dispersed gapless photon mode.

To understand the meaning of the gapless mode, we note that due to the non-zero Hall conductances ± 1 for the spin-up and spin-down spinons, a flux quantum of the gauge field a_μ will induce one spin-up spinon and minus one spin-down spinon. Thus a flux quantum carries a spin quantum number $S_z = 1$. The "magnetic field" of a_μ is nothing but the S_z density. Therefore, the linear gapless mode of a_μ is actually the density fluctuations of S_z. The appearance of a linear gapless mode of a conserved density as the only low lying excitation implies that the corresponding symmetry is spontaneously broken in the ground state.[14, 15] Thus the LL state contains an XY order: $\langle S^+\rangle \neq 0$ and the gapless gauge field a_μ is the Goldstone boson mode of the $U(1)$ symmetry breaking. The "electric charge" of a_μ gauge field corresponds to the vortex in the XY ordered phase. This duality [16, 17] between the spin degrees of freedom and the gauge degrees of freedom allows us to translate the physics in the two languages back and forth.

For example let us consider the monopole insertion operator $V^\dagger$ which inserts a 2π gauge flux. $V^\dagger$ flips one spin as mentioned before and can be expanded as $V^\dagger \sim \sum_i e^{i\theta_i} S_i^+$, where the summation is restricted to the area where the 2π flux is inserted. Because $V^\dagger$ has long-range correlation in the Coulomb phase, by the above duality, the spin correlation $\langle e^{i\theta_i} S_i^+, e^{-i\theta_j} S_j^-\rangle$ is also long ranged.

To study what kind of the XY spin order that the LL state has, we have calculated spin-spin correlation function $\langle S_{x,i}S_{x,j} + S_{y,i}S_{y,j}\rangle$ of the projected LL state and found the XY ordered pattern as shown in Fig.2(b). This order pattern is referred as the $q = 0$ magnetic order in literatures[18]. To confirm this is indeed the order pattern we directly compute the relative phases of S_i^+ operators in the monopole insertion operator $V^\dagger$:

$$\theta_j - \theta_i = \arg\Big(\frac{\langle n+1 \text{ flux quanta}|PS_j^+P|n \text{ flux quanta}\rangle}{\langle n+1 \text{ flux quanta}|PS_i^+P|n \text{ flux quanta}\rangle}\Big), \tag{7}$$

where $|n \text{ flux quanta}\rangle$ is a LL state with n flux quanta and P is the projection operator. To understand this formula one should note that $P|n+1 \text{ flux quanta}\rangle = V^\dagger P|n \text{ flux quanta}\rangle$, where $V^\dagger$ inserts an extra 2π flux uniformly while flipping one spin, and thus $V^\dagger \sim \sum_i e^{i\theta_i} S_i^+$ with summation over all sites. These relative phases can be computed analytically by choosing proper gauge (i.e., gauge that respects physical symmetries). To get the answer quickly we perform the Monte Carlo calculation on a finite lattice. For a 6 by 6 unit cell sample on torus with $n = 4$, we compute this relative phase for 9 sites within the three adjacent unit cells and the result is given in Fig. 2(b). We find the relative phases are $0, \pm\frac{2\pi}{3}$ alternating between the three sublattices and this is exactly the 120° $q = 0$ order pattern.[26]

Because $V^\dagger$ is in fact the magnetization operator, we conclude that the in-plane staggered magnetization $M \sim B^\alpha$ scales with the external magnetic field, where the exponent α is the scaling dimension of the monopole operator at the Dirac-SL fixed point, because B is dimension-1. This exponent is computable by numerical simulations of QED$_3$ or the field theory techniques such as $1/N$ expansions, and subject to further study.

So far we show that the FP state is unstable towards the S_z-broken LL state by numerical arguments. In the following we present an analytical argument that the S_z symmetry is broken by studying the low energy effective theory. The FP state is characterized by a electron-like

Fermi pocket of spin up spinons $f_{k\uparrow}$ and a hole-like Fermi pocket of spin down spinons $f_{k\downarrow}$. After doing a particle-hole transformation on the spin down Fermi pocket only $f_{k\downarrow} \to h_{-k\uparrow}$ we have $f_{k\uparrow}$ and $h_{-k\uparrow}$ carry same spin but opposite gauge charges. The "Coulomb" attraction between the two particles will cause a pairing instability $\langle f^\dagger_{k\uparrow} h^\dagger_{-k\uparrow}\rangle = \langle f^\dagger_{k\uparrow} f_{k\downarrow}\rangle \neq 0$. This is a triplet excitonic insulator and this has been discussed in the context of graphene[19, 20]. Besides Coulomb attraction, there is also an Amperean attraction[21] between the currents of $f^\dagger_{k\uparrow}$ and $h^\dagger_{-k\uparrow}$ excitations. The "Coulomb" attraction and the Amperean attraction are cooperating for the same condensation. Note that $f^\dagger_{k\uparrow} f_{k\downarrow}$ is a S_z spin-1 object.

Finally we discuss the consequences of this spontaneous spin ordering in experiment. First the in-plane $q = 0$ magnetization pattern and its scaling law $M \sim B^\alpha$ is observable by neutron scattering. Second since the ground state breaks the parity and S_z rotation symmetry, it is separated from the high temperature paramagnetic phase by at least one *finite* temperature phase transition. This transition can be first order or continuous. If the transition is a continuous one that restores the S_z-$U(1)$ symmetry, we expect it to be the Kosterlitz-Thouless universality. Here we simply estimate the transition temperature $T_c \sim \frac{\mu_B B}{k_B}$ when $\mu_B B \lesssim \chi^{Prj}$. This is because the external magnetic field B is the only energy scale if it is much smaller than the spinon band width χ^{Prj}. Since these are intrinsic properties of the Dirac-SL in magnetic field, they can be used to experimentally detect the possible Dirac-SL ground state in Herbertsmithite and other materials where a Dirac-SL may be realized. This XY spin ordering can also serve a way to detect Dirac-SL in numerical studies of the Kagome lattice Heisenberg model such as exact diagonalization.

There are other proposals for the possible non-magnetic Valence Bond Solid (VBS) ground states in the spin-1/2 Heisenberg model on Kagome lattice[22, 23]. If one of those VBS states is the ground state of Herbertsmithite, it will surely not break S_z symmetry in a small magnetic field because the VBS phases are fully gapped. The S_z symmetry can be broken at an external magnetic field larger than the spin gap due to the triplon condensation. However the XY magnetic order generated in this fashion is unlikely to be the $q = 0$ pattern because the VBS orders itself breaks translation and we expect their XY orders also breaks translation and have a large unit cell. Therefore the spontaneous spin ordering we present in this work can be used to differentiate the $U(1)$-Dirac SL from VBS states experimentally.

It is likely that in the current Herbertsmithite compound there is an energy scale below which the significant amount of impurities and/or the Dzyaloshinskii-Moriya(DM) interaction start to play an important role. Recent experiments estimate the strength of the DM interaction to be $\sim$ 15K[24]. In a strong magnetic field $\sim$ 30 tesla we expect that it is possible to suppress their effects and reveal the intrinsic property of the Dirac-SL.

We thank M. Hermele, T. Senthil and F.-C. Zhang for helpful discussions and T. M. Rice for raising the issues concerning graphene. This research is supported by NSF grant DMR-0706078 and DMR-0804040.

---

[1] O. Ofer, A. Keren, E. A. Nytko, M. P. Shores, B. M. Bartlett, D. G. Nocera, C. Baines, and A. Amato (2006), URL `cond-mat/0610540`.

[2] J. S. Helton, K. Matan, M. P. Shores, E. A. Nytko, B. M. Bartlett, Y. Yoshida, Y. Takano, A. Suslov, Y. Qiu, J. H. Chung, et al., Phys. Rev. Lett. **98**, 107204 (2007).

[3] P. Mendels, F. Bert, M. A. de Vries, A. Olariu, A. Harrison, F. Duc, J. C. Trombe, J. Lord, A. Amato, and C. Baines, Phys. Rev. Lett. **98**, 077204 (2007).

[4] G. Misguich and P. Sindzingre, Eur. Phys. J. B **59**, 305 (2007).

[5] M. Rigol and R. R. P. Singh, Phys. Rev. B **76**, 184403 (2007).

[6] Y. Ran, M. Hermele, P. A. Lee, and X.-G. Wen, Phys. Rev. Lett. **98**, 117205 (2007).

[7] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, Nature **438**, 197 (2005).

[8] Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, Nature **438**, 201 (2005).

[9] A. N. Redlich, Phys. Rev. D **29**, 2366 (1984).

[10] J. B. Marston, Phys. Rev. Lett. **64**, 1166 (1990).

[11] C. Gros, Annals of Physics **189**, 53 (1989).

[12] M. B. Hastings, Phys. Rev. B **63**, 014413 (2000).

[13] M. Hermele, Y. Ran, P. A. Lee, and X.-G. Wen, Phys. Rev. B **77**, 224413 (2008).

[14] X. G. Wen and A. Zee, Phys. Rev. B **41**, 240 (1990).

[15] X.-G. Wen and A. Zee, Phys. Rev. Lett. **69**, 1811 (1992).

[16] C. Dasgupta and B. I. Halperin, Phys. Rev. Lett. **47**, 1556 (1981).

[17] M. P. A. Fisher and D. H. Lee, Phys. Rev. B **39**, 2756 (1989).

[18] A. B. Harris, C. Kallin, and A. J. Berlinsky, Phys. Rev. B **45**, 2899 (1992).

[19] D. V. Khveshchenko and W. F. Shively, Phys. Rev. B **73**, 115104 (2006).

[20] I. L. Aleiner, D. E. Kharzeev, and A. M. Tsvelik, Phys. Rev. B **76**, 195415 (2007).

[21] S.-S. Lee, P. A. Lee, and T. Senthil, Phys. Rev. Lett. **98**, 067006 (2007).

[22] P. Nikolic and T. Senthil, Phys. Rev. B **68**, 214415 (2003).

[23] R. R. P. Singh and D. A. Huse, Phys. Rev. B **76**, 180407 (2007).

[24] A. Zorko, S. Nellutla, J. van Tol, L. C. Brunel, F. Bert, F. Duc, J.-C. Trombe, M. A. de Vries, A. Harrison, and P. Mendels, Phys. Rev. Lett. **101**, 026405 (2008).

[25] Y. Ran, A. Vishwanath, and D.-H. Lee (2008), URL `arXiv.org:0806.2321`.

[26] We actually find that this relative phase is independent of n. Even when $n = 0$, namely the pure $U(1)$-Dirac SL state, the relative phase is the same as the $n = 4$ data that we presented up to statistical error. This indicates that the relative phase which we computed is an

intrinsic property of the $U(1)$-Dirac SL, i.e., they are the spin-1 monopole quantum numbers. The Eq.(7) is a systematic way to compute monopole quantum numbers in any $U(1)$-Dirac SL. A more detailed discussion on this method is presented as an appendix in Ref.[25]. The monopole quantum numbers obtained in this method (see Fig. 2(b)) are consistent with the results obtained by an independent study on the Dirac SL in zero magnetic field based on symmetry group analysis[13].